



# TSG-6 Regulates Bone Remodeling through Inhibition of Osteoblastogenesis and Osteoclast Activation

Submitted by Emmanuel Lemoine on Tue, 06/10/2014 - 11:21

Titre	TSG-6 Regulates Bone Remodeling through Inhibition of Osteoblastogenesis and Osteoclast Activation
Type de publication	Article de revue
Auteur	Mahoney, David-J. [1], Mikecz, Katalin [2], Ali, Tariq [3], Mabilletau, Guillaume [4], Benayahu, Dafna [5], Plaas, Anna [6], Milner, Caroline-M. [7], Day, Anthony-J. [8], Sabokbar, Afsie [9]
Editeur	American Society for Biochemistry and Molecular Biology
Type	Article scientifique dans une revue à comité de lecture
Année	2008
Langue	Anglais
Date	2008/09/19
Numéro	38
Pagination	25952 - 25962
Volume	283
Titre de la revue	Journal of Biological Chemistry
ISSN	0021-9258, 1083-351X
Résumé en anglais	<p>TSG-6 is an inflammation-induced protein that is produced at pathological sites, including arthritic joints. In animal models of arthritis, TSG-6 protects against joint damage; this has been attributed to its inhibitory effects on neutrophil migration and plasmin activity. Here we investigated whether TSG-6 can directly influence bone erosion. Our data reveal that TSG-6 inhibits RANKL-induced osteoclast differentiation/activation from human and murine precursor cells, where elevated dentine erosion by osteoclasts derived from TSG-6<sup>-/-</sup> mice is consistent with the very severe arthritis seen in these animals. However, the long bones from unchallenged TSG-6<sup>-/-</sup> mice were found to have higher trabecular mass than controls, suggesting that in the absence of inflammation TSG-6 has a role in bone homeostasis; we have detected expression of the TSG-6 protein in the bone marrow of unchallenged wild type mice. Furthermore, we have observed that TSG-6 can inhibit bone morphogenetic protein-2 (BMP-2)-mediated osteoblast differentiation. Interaction analysis revealed that TSG-6 binds directly to RANKL and to BMP-2 (as well as other osteogenic BMPs but not BMP-3) via composite surfaces involving its Link and CUB modules. Consistent with this, the full-length protein is required for maximal inhibition of osteoblast differentiation and osteoclast activation, although the isolated Link module retains significant activity in the latter case. We hypothesize that TSG-6 has dual roles in bone remodeling; one protective, where it inhibits RANKL-induced bone erosion in inflammatory diseases such as arthritis, and the other homeostatic, where its interactions with BMP-2 and RANKL help to balance mineralization by osteoblasts and bone resorption by osteoclasts.</p>
URL de la notice	<a href="http://okina.univ-angers.fr/publications/ua3290">http://okina.univ-angers.fr/publications/ua3290</a> [10]

DOI 10.1074/jbc.M802138200 [11]  
Lien vers le document <http://dx.doi.org/10.1074/jbc.M802138200> [11]

---

### Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=4576](http://okina.univ-angers.fr/publications?f[author]=4576)
- [2] [http://okina.univ-angers.fr/publications?f\[author\]=4577](http://okina.univ-angers.fr/publications?f[author]=4577)
- [3] [http://okina.univ-angers.fr/publications?f\[author\]=4578](http://okina.univ-angers.fr/publications?f[author]=4578)
- [4] <http://okina.univ-angers.fr/guillaume.mabilleau/publications>
- [5] [http://okina.univ-angers.fr/publications?f\[author\]=4579](http://okina.univ-angers.fr/publications?f[author]=4579)
- [6] [http://okina.univ-angers.fr/publications?f\[author\]=4580](http://okina.univ-angers.fr/publications?f[author]=4580)
- [7] [http://okina.univ-angers.fr/publications?f\[author\]=4581](http://okina.univ-angers.fr/publications?f[author]=4581)
- [8] [http://okina.univ-angers.fr/publications?f\[author\]=4582](http://okina.univ-angers.fr/publications?f[author]=4582)
- [9] [http://okina.univ-angers.fr/publications?f\[author\]=4570](http://okina.univ-angers.fr/publications?f[author]=4570)
- [10] <http://okina.univ-angers.fr/publications/ua3290>
- [11] <http://dx.doi.org/10.1074/jbc.M802138200>

Publié sur *Okina* (<http://okina.univ-angers.fr>)